

Using Virtual Reality to elicit Empathy: a narrative review

Emilija Kizhevska*

emilija.kizhevska@ijs.si

Jožef Stefan Institute, Jožef Stefan International

Postgraduate School (IPS)

Ljubljana, Slovenia

Tiago Guerreiro

LASIGE, Faculdade de Ciências da Universidade de Lisboa

Lisbon, Portugal

tjvg@di.fc.ul.pt

Filipa Ferreira-Brito

LASIGE, Faculdade de Ciências da Universidade de Lisboa,
Portugal;

ISAMB, Faculdade de Medicina da Universidade de Lisboa
Lisbon, Portugal

filipa.r.brito@gmail.com

Mitja Luštrek

Jožef Stefan Institute, Jožef Stefan International

Postgraduate School (IPS)

Ljubljana, Slovenia

mitja.lustrek@ijs.si

ABSTRACT

Virtual reality (VR) technology is often characterized as “the ultimate empathy machine” as it enables users to experience how it is to be someone else or be somewhere other than where they are in the real physical world. Here, we conducted a narrative review of studies focused on using VR to elicit empathy. Considering the synthesized literature, we identified three contexts where VR systems have been used as a tool to study empathic behavior, namely: 1) to promote pro-environmental behavior; 2) to promote prosocial behavior toward specific social groups (e.g., refugees); and 3) to medical training to promote empathy and more in-depth knowledge of clinical conditions. Based on the data collected VR seems more effective in evoking empathy than traditional approaches such as films/videos, narratives, and curriculum content. Furthermore, it was possible to identify an increase in participants’ empathic responses immediately after exposure to VR and up to some period after the intervention (e.g., two months). However, despite the popularity of VR in the study of empathy, the conclusions that can be drawn regarding VR efficacy to promote/ elicit empathic behavior are still obscured by the lack of consensual theoretical constructs, the use of a wide variety of self-reported measures, and the incipient use of physiological measures.

KEYWORDS

virtual reality, empathy, 360° immersive virtual environment

ACM Reference Format:

Emilija Kizhevska, Filipa Ferreira-Brito, Tiago Guerreiro, and Mitja Luštrek. 2022. Using Virtual Reality to elicit Empathy: a narrative review. In *Proceedings of Workshop on Virtual Reality for Health and Wellbeing (MUM) Workshop on VR for Health and Wellbeing @ MUM 2022, Lisbon, Portugal (November, 2022)*, 4 pages.

Permission to make digital or hard copies of all or part of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for components of this work owned by others than ACM must be honored. Abstracting with credit is permitted. To copy otherwise, or republish, to post on servers or to redistribute to lists, requires prior specific permission and/or a fee. Request permissions from permissions@acm.org.

MUM, November 27–30, 2022, Lisbon, Portugal

© 2022 Association for Computing Machinery.

ACM ISBN 978-x-xxxx-xxxx-x/YY/MM...\$15.00

<https://doi.org/XXXXXXXXXXXXXX>

1 INTRODUCTION

Virtual reality (VR) can be defined as an immersive and interactive computer-generated environment that gives users the feeling of being somewhere else other than where they are in the physical world [2]. Through VR, individuals have the feeling of being in a real environment and, potentially, behave accordingly [12]. Thereby, a considerable number of scientific papers have presented claims that VR can be used to elicit empathic behavior, characterizing VR technology as “the ultimate empathy machine” [3]. Empathy is the ability or tendency to share and understand others’ thoughts, emotions, and internal states. Although there is no widely accepted definition of empathy, there is a consensus regarding its multidimensional nature. For instance, one of the most used theoretical models is the one that conceives empathy as encompassing a cognitive component (i.e., the capacity for understanding another person’s experience and perspective) and an emotional component (i.e., the ability to share the emotional state of another person) [8]. Furthermore, in the last few years, emotion regulation strategies have shown that empathy with other emotional competencies such as mindfulness, self-compassion, and resilience are predictors of well-being [10]. Recent studies have concluded when empathy declines, distress is a key determinant of it [11]. Furthermore, empathy has been linked to increased well-being, reduced symptoms of burnout, and more meaningful work experiences as in the case of medical workers [14]. This narrative review is a first attempt to map the current state-of-the-art on VR to study empathic behavior since, to the best of our knowledge, there is no review on this topic without focusing on a specific context (e.g., medical training) or populations (e.g., schizophrenia).

2 METHODS

We conducted a narrative review of articles that use VR technology to study human empathic responses. Papers were screened independently by two researchers (EK and FFB), and data extraction was performed by one of the co-authors (EK) and checked/complemented by another co-author (FFB).

2.1 Electronic Databases and Search String

For the database search, we used a convenience pool of databases (i.e., the ones that authors are more familiar with). PubMed/MEDLINE

and EBSCO (Psychology and Behavioral Sciences Collection) electronic databases were searched on the 14th of September using the following search string:

((empathy OR (“cognitive empathy”) OR (“emotional empathy”)) AND (“Virtual Reality”) OR (VR)))

However, we plan to increase the number of databases include in the next step of our work which will be conducting a systematic review of VR to study empathy.

2.2 Filters and Eligibility criteria

Whenever possible, electronic searches were restricted to full-text papers published in peer-reviewed journals between 2012 – 2022 in English. Furthermore, in the EBSCO electronic database, papers were searched using the broader search field, while in PubMed/MEDLINE, the search-string was searched using the title/abstract search field. All papers retrieved were assessed against the following inclusion criteria: 1) aimed to study participants’ empathic behavior [even if the authors did not directly mention the term empathy]; 2) while or after being exposed to VR. Book chapters, gray literature, and systematic and narrative reviews were excluded. Furthermore, papers were excluded if they recruited exclusively under-age participants (i.e., < 18 years) or did not present standardized measures of empathy (e.g., feasibility studies).

3 RESULTS

A total of 42 papers were retrieved from database searches and exported to the Rayyan QCRI web application [13]. Papers were assessed against the eligibility criteria as depicted in Fig1.




Figure 1: Flow diagram of the studies included

3.1 General characteristics of the studies included

The potential of VR to elicit empathy was analyzed in three different contexts: 1) as a means to promote pro-environmental behaviors, for instance, by increasing donations to protect coral reefs [15]; 2) to promote empathy and more in-depth knowledge of the difficulties faced by patients with dementia in healthcare workers [9] and caregivers [4]; and 3) to promote prosocial behavior by facilitating perspective taking on a peer [1], homeless people [5], refugees [18], victims of sexual harassment [17], or towards people/avatars experiencing pain [16] and pleasure [6][7]. A total of 2061 participants were recruited with sample sizes ranging between 24 [6][18] and 1006 [15] participants, with four studies having recruited more than 100 participants (114 [9], 180 [1], 556 [5], and 1006 [15] participants, respectively). On average, participants were 33.19 +/- 12.75 years [19.91 - 55.1] (mean age), and 56.85% identified themselves as female, with one study only recruiting men [17]. Regarding exposure to VR, sessions took on average 10.18 +/- 5.34 minutes [4.87 - 20] (mean min), with six studies [15][5][7][6][4][18] immersed participants in one session and three immersed participants twice [1][17][16].

3.2 Measures of Empathy and VR features analyzed

With no exceptions, all included papers reported using one [15][9][7][6][4][18] or multi-self-reported questionnaires [1][5][17][16] to measure empathy. Examples of the questionnaires used are the Interpersonal Reactivity Index [5][6], or some of its sub-scales [9][7], the Empathy Scale [17], the Situational Empathy and Perspective Taking Scale [9], the Empathy for Pain Scale [7][6], or adapted items from other empathy scales [4][18]. In addition, three papers also assessed the impact of VR on the feeling of connectedness with others [17][5][1], body transfer [1], attitude towards gender-based violence [17], personal distress, attitudes towards the homeless [5] and social presence [5][16]. Regarding objective measures, four studies used biometric signals, such as galvanic skin response (GSR) [16], and skin conductance reactivity alone (SCR) [7][6] or combined with heart rate (HR) [6]. Also, participants’ head position was tracked by a wireless InterSense IS-900 VET system [16] or using an infrared camera (Oculus DK2 IR camera) [5]. Concerning the analysis of which VR/technical components were more effective to elicit empathy, studies compared the impact of 360° immersive virtual environments (IVE) with two-dimensional video/film [15][18], curriculum contents (i.e., workshops [9] or e-courses material [4]), narrative-based perspective taking exercises [5][17] text-based information [15][5]. Other technical features analyzed were the degree of immersion (i.e., 3D versus 2D) [5] and the perspective in which participants experienced the VE (i.e., first-person versus third-person perspective) [7][6]. The majority of the papers immersed participants using head-mounted displays (HMD) [17], some have specified it is Oculus Rift DK2 [5][7][6][4] or HTC Vive [1], but also headsets as Oculus Quest [9], Samsung VR headset [18] or Zeiss VR One Headset [15] were used, and in one study the immersion is achieved by 3D shutter glasses NuVision [16].

3.3 Efficacy of VR to elicit empathy

The impact of VR to elicit empathy seems to be moderated by several factors (e.g., contexts, order of exposure, control condition to which VR is compared, and demographic variables). For instance, two studies reported that 360° IVEs are more effective than (unidirectional) videos/films to promote pro-environmental behavior [15] and empathy toward refugees [18]. In another study, targeting personal distress and empathy toward the homeless [5], no statistical differences were found between 360° IVEs and narrative-based perspective exercises. However, it was possible to identify a long-lasting positive attitude toward the homeless (i.e., two months after the intervention) in the IVEs group but not in the control group [5]. In addition, when analyzing empathy toward victims of sexual harassment, 360° IVEs was as effective as a narrative-based perspective-taking exercise [17]. However, significantly higher scores on empathy were identified when participants were first exposed to the narrative-based perspective-taking exercise after the 360° IVEs rather than before the 360° video [17]. When VR was used as a medical training tool, one study [4] showed that 360° IVEs are more effective in promoting empathy toward the challenges faced by people with dementia, while in another study, 360° IVEs' impact on medical training and empathy towards patients with dementia were moderated by demographic characteristics, with significant improvements on empathy levels being identified only in older and non-english native speakers participants [9]. Furthermore, higher scores on empathy scales were identified when participants are asked to interact with avatars depicting familiar faces expressed pain compared to unfamiliar avatars [16] or when taking the perspective of an individual they expect later to interact [1]. Finally, in the studies where participants watch/embody people/avatars experiencing pain and pleasure it was found higher feelings of ownership in the first-person perspective compared to the third-person perspective condition [7][6].

4 CONCLUSION

Our review provides an overview of the current state of research concerning the use of VR to study/ elicit empathy. As the next step, we plan to upgrade it to a systematic review by surveying more papers and analyzing them more thoroughly. While we find the evidence on the effectiveness of VR in eliciting empathy promising, it is still clouded by the lack of gold-standard instruments to measure empathy. We believe that an important next research step in this domain should be finding how empathy influences physiological attributes and using the sensors as a more objective measure of empathy. Furthermore, VR's impact on empathy seems to be moderated by demographic, methodological and context variables that should be taken into consideration in future studies, too.

ACKNOWLEDGMENTS

The research and the preparation of the paper on the part of E.K. was supported by the Slovenian Research Agency (ARRS) as part of the young researcher PhD program, Grant (PR-12092). FCT through LASIGE funding (ref. UIDB/00408/2020 and ref. UIDP/00408/2020). The collaboration is supported and funded by the WideHealth project. The WideHealth project has received funding from the

European Union's Horizon 2020 research and innovation program under grant agreement No 952279.

REFERENCES

(*Asterisk denotes studies included in the review)

- [1] *Van Loon A, Bailenson J, Zaki J, Bostick J, and Willer R. 2018. Virtual reality perspective-taking increases cognitive empathy for specific others. *Plos one* 13, 8 (8 2018).
- [2] Sun Joo (Grace) Ahn, Joshua Bostick, Kristine L. Nowak Elise Ogle, Kara T. McGillicuddy, and Jeremy N. Bailenson. 2016. Experiencing Nature: Embodying Animals in Immersive Virtual Environments Increases Inclusion of Nature in Self and Involvement with Nature. *Journal of Computer-Mediated Communication* 21, 6 (Nov. 2016), 399–419. <https://doi.org/10.1111/jcc4.12173>
- [3] Barbot, Baptiste, and James C. Kaufman. 2020. What makes immersive virtual reality the ultimate empathy machine? Discerning the underlying mechanisms of change. *Computers in Human Behavior* 111, 106431 (Oct. 2020).
- [4] *Wijma EM, Veerbeek MA, Prins M, Pot AM, and Willemse BM. 2018. A virtual reality intervention to improve the understanding and empathy for people with dementia in informal caregivers: results of a pilot study. *Aging mental health* 22, 9 (Sept. 2018), 1121–9.
- [5] *Herrera F, Bailenson J, Weisz E, Ogle E, and Zaki J. 2018. Building long-term empathy: A large-scale comparison of traditional and virtual reality perspective-taking. *Plos one* 13, 10 (Oct. 2018).
- [6] *Fusaro, Martina, Gaetano Tieri, and Salvatore Maria Aglioti. 2016. Seeing pain and pleasure on self and others: behavioral and psychophysiological reactivity in immersive virtual reality. *Journal of Neurophysiology* 116, 6 (Dec. 2016), 2656–2662.
- [7] *M. Fusaro, G. Tieri, and S. M. Aglioti. 2019. Influence of cognitive stance and physical perspective on subjective and autonomic reactivity to observed pain and pleasure: An immersive virtual reality study. *Consciousness and cognition* 67, 67 (Jan. 2019), 86–97.
- [8] Harari H, Shamay-Tsoory SG, Ravid M, and Levkovitz Y. 2010. Double dissociation between cognitive and affective empathy in borderline personality disorder. *Psychiatry research* 175, 3 (Feb. 2010), 277–9.
- [9] *Stargatt J, Bhar S, Petrovich T, Bhowmik J, Sykes D, and Burns K. 2021. The effects of virtual reality-based education on empathy and understanding of the physical environment for dementia care workers in Australia: A controlled study. *Journal of Alzheimer's Disease* 84, 3 (Jan. 2021), 1247–57.
- [10] Zwack J and Schweitzer J. 2013. If every fifth physician is affected by burnout, what about the other four? Resilience strategies of experienced physicians. *Academic Medicine* 88, 3 (March 2013), 382–9.
- [11] Neumann M, Edelhäuser F, Tauschel D, Fischer MR, Wirtz M, Woopen C, Haramati A, and Scheffer C. 2011. Empathy decline and its reasons: a systematic review of studies with medical students and residents. *Academic Medicine* 86, 8 (Aug. 2011), 996–1009.
- [12] Sanchez-Vives M. and Slater M. 2005. From presence to consciousness through virtual reality. *Nat Rev Neurosci* 6, 4 (April 2005), 332–339. <https://doi.org/10.1038/nrn1651>
- [13] Mourad Ouzzani, and Hossam Hammady, and Zbys Fedorowicz, and Ahmed Elmagarmid. 2016. Rayyan — a web and mobile app for systematic reviews. *Systematic Reviews*, 5:210. 10.1186/s13643-016-0384-4
- [14] Thomas MR, Dyrbye LN, Huntington JL, Lawson KL, Novotny PJ, Sloan JA, and Shanafelt TD. 2007. How do distress and well-being relate to medical student empathy? A multicenter study. *Journal of general internal medicine* 22, 2 (Feb. 2007), 177–83.
- [15] *Nelson, Katherine M., Eva Anggraini, and Achim Schlüter. 2020. Virtual reality as a tool for environmental conservation and fundraising. *Plos one* 15, 4 (April 2020).
- [16] *Bouchard S, Bernier F, Boivin É, Dumoulin S, Laforest M, Guitard T, Robillard G, Monthuy-Blanc J, and Renaud P. 2013. Empathy toward virtual humans depicting a known or unknown person expressing pain. *Cyberpsychol Behav Soc Netw* 16, 1 (Jan. 2013), 61–71.
- [17] *Ventura S, Cardenas G, Miragall M, Riva G, and Baños R. 2021. How does it feel to be a woman victim of sexual harassment? The effect of 360°-video-based virtual reality on empathy and related variables. *Cyberpsychology, Behavior, and Social Networking* 24, 4 (April 2021), 258–66.
- [18] *Schutte, Nicola S., and Emma J. Stilianović. 2017. Facilitating empathy through virtual reality. *Motivation and emotion* 41, 6 (Dec. 2017), 708–712.